

TYPES of Myositis

do not have skin rashes. Some PM patients have trouble swallowing (dysphagia), difficulty breathing (often associated with interstitial lung disease) and muscle pain. PM may be associated with a malignancy or with other autoimmune diseases.

Inclusion-Body Myositis (IBM)

is found in more men than women and is rarely seen in people less than 50 years old. IBM progresses more slowly than the other types of myositis, as weakness happens gradually over months or years. Some of the first signs of IBM are falling, difficulty standing from a seated position and weakening grip. Muscles most often affected are those at the front of the thighs, hips, foot-elevators, fingers, wrists, upper arms, shoulders, neck and back. Many IBM patients notice shrinking, or atrophy, in the arms and thighs as the muscles become weaker. Trouble swallowing (dysphagia) is a common problem for IBM patients. Weakness of facial muscles is sometimes seen.

Juvenile Forms of Myositis (JM)

occur in children 18-years-old and younger, and affect more girls than boys. Juvenile dermatomyositis (JDM) is the most common form, affecting children in Canada. Polymyositis in children (JPM) occurs in



*Juvenile Forms of Myositis
found in children
18-years-old and younger*

TYPES of Myositis

approximately 10% of children with JM. Signs of JM include the characteristic skin rashes (i.e. a visible, reddish-purple rash over the eyelids or over joints), trouble climbing or lifting the head, weak voice (dysphonia) or problems swallowing (dysphagia). The muscles most affected are those closest to the centre of the body – neck, stomach, upper arms and legs. About half of the children with JM report pain in their muscles. Some children have calcinosis (hardened lumps under the skin) or contractures (when muscle shortens and causes joints to stay bent). Children may have one or more other autoimmune disease(s) along with myositis.

#GiveMeMuscle
www.myositis.ca



#103, 7724 Bowness Road NW
Calgary, Alberta, T3B 0H1



MYOSITIS
CANADA

*Supporting Community in
Myositis Diagnosis, Treatment and
Rehabilitation*

WHO we are

Myositis Canada is a non-profit, voluntary agency dedicated to improving the lives of people affected by myositis. The agency was formed in 2015 by people with myositis.

We are governed by a volunteer Board of Directors that includes patients, family members, and interested professionals, bringing diverse strengths and perspectives to the organization.

Myositis Canada envisions a world where each new case of myositis is diagnosed and cured within months, and where full rehabilitation from myositis damage is routine.



Brenda is a Canadian
"Face" of Myositis

Myositis Canada believes in integrity, honesty, teamwork, community and most importantly, respect.

OUR Mission

To support people with myositis, caregivers, researchers, and health care providers, in the diagnosis and treatment of, and rehabilitation from myositis.

WHAT is Myositis?

Myositis describes inflammation or swelling of the muscle tissue. General muscle inflammation can occur after exercising or taking certain medication, or it can be from one of the chronic inflammatory muscle disorders.

Dermatomyositis (*DM*), polymyositis (*PM*), inclusion-body myositis (*IBM*), and juvenile forms of myositis (*JM*) are all inflammatory myopathies or diseases where there is inflammation and loss of muscle. **Depending on the type of myositis, the first signs of myositis may include general fatigue, trouble standing from a seated position, difficulty climbing stairs and weak grasping of objects.**

The underlying causes of DM, PM, IBM and JM are not known, but some doctors believe there is an environmental exposure (perhaps to an infection or sunlight) that triggers the disease in someone who has certain specific, but not yet fully defined genes or gene sequences that predispose him or her. Inflammatory myopathies are classified as autoimmune diseases, meaning the body's immune system, which normally fights infections and virus, does not stop fighting once the infection or virus is gone. **The immune system is misdirected and attacks the body's own normal, healthy tissue.** Inflammatory myopathies are rare diseases and combined, affect an estimated 50,000 people in Canada. Another 2,500 people are likely to have myositis but are misdiagnosed.

TYPES of Myositis

Dermatomyositis (DM)

affects people of any age or sex but is more common in women than men. DM is the easiest type of myositis to diagnose because it typically has a visible skin rash caused by inflammation of blood vessels under the skin. The DM rash looks patchy and reddish or purple, and is found on the eyelids, elbows, knees and knuckles. Additional rashes may occur on the cheeks, nose, back and upper chest. Some people also have hardened bumps under the skin, called calcinosis. Patients usually experience gradual muscle weakness and sometimes pain, and they often report the rash well before the muscle weakness. Three sub-types of DM are amyopathic DM, where the skin is affected but muscles are not involved; cancer-associated DM, where cancer and dermatomyositis are diagnosed within two to three years of one another; and overlap, where DM is associated with other autoimmune diseases.

Polymyositis (PM)

is found more frequently in adults than children. More women than men have PM. Patients experience muscle weakness gradually over weeks or months, typically beginning with muscles closest to and within the trunk of the body, such as neck, hip, back, and shoulder muscles, and affecting both sides of the body equally. Some patients also have weakness in their hands and fingers. PM patients generally